1: NP_443204. Reports leucine-rich alph...[gi:16418467]

BLink, Conserved Domains, Links

Comment Features Sequence

```
LOCUS       NP_443204                347 aa            linear   PRI 25-OCT-2001
DEFINITION  leucine-rich alpha-2-glycoprotein [Homo sapiens].
ACCESSION   NP_443204
VERSION     NP_443204.1  GI:16418467
DBSOURCE    REFSEQ: accession NM_052972.1
KEYWORDS    .
SOURCE      Homo sapiens (human)
  ORGANISM  Homo sapiens
            Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
            Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE   1  (sites)
  AUTHORS   Bunkenborg,J., Pilch,B.J., Podtelejnikov,A.V. and Wisniewski,J.R.
  TITLE     Screening for N-glycosylated proteins by liquid chromatography mass
            spectrometry
  JOURNAL   Proteomics 4 (2), 454-465 (2004)
   PUBMED   14760718
REFERENCE   2  (residues 1 to 347)
  AUTHORS   Takahashi,N., Takahashi,Y. and Putnam,F.W.
  TITLE     Periodicity of leucine and tandem repetition of a 24-amino acid
            segment in the primary structure of leucine-rich alpha
            2-glycoprotein of human serum
  JOURNAL   Proc. Natl. Acad. Sci. U.S.A. 82 (7), 1906-1910 (1985)
   PUBMED   3856868
REFERENCE   3  (residues 1 to 347)
  AUTHORS   O'Donnell,L.C., Druhan,L.J. and Avalos,B.R.
  TITLE     Molecular characterization and expression anaylysis of leucine-rich
            alpha-2-glycoprotein, a novel marker of granulocytic
            differentiation
  JOURNAL   Unpublished
COMMENT     PROVISIONAL REFSEQ: This record has not yet been subject to final
            NCBI review. The reference sequence was derived from AF403428.1.
FEATURES             Location/Qualifiers
     source          1..347
                     /organism="Homo sapiens"
                     /db_xref="taxon:9606"
                     /chromosome="19"
                     /map="19p13.3"
                     /clone="IMAGE:2403704"
     Protein         1..347
                     /product="leucine-rich alpha-2-glycoprotein"
                     /calculated_mol_wt=38178
     sig_peptide     1..347
                     /note="putative"
                     /calculated_mol_wt=38178
```

```
     Region          <80..241
                     /region_name="LRR_RI"
                     /note="Leucine-rich repeats (LRRs), ribonuclease inhibitor
                     (RI)-like subfamily. LRRs are 20-29 residue sequence
                     motifs present in many proteins that participate in
                     protein-protein interactions and have different functions
                     and cellular locations; cd00116"
                     /db_xref="CDD:29015"
     Site            186
                     /site_type="glycosylation"
                     /experiment="experimental evidence, no additional details
                     recorded"
                     /citation=[1]
     Region          299..346
                     /region_name="LRRCT"
                     /note="Leucine rich repeat C-terminal domain; smart00082"
                     /db_xref="CDD:47428"
     Region          299..346
                     /region_name="Leucine rich repeat C-terminal domain"
                     /note="LRRCT; Leucine rich repeat C-terminal domain"
                     /db_xref="CDD:smart00082"
     Region          299..346
                     /region_name="Leucine rich repeat C-terminal domain.
                     Leucine Rich Repeats pfam00560 are short sequence motifs
                     present in a number of proteins with diverse functions and
                     cellular locations. Leucine Rich Repeats are often flanked
                     by cysteine rich domains. This domain is often found at
                     the C-terminus of tandem leucine rich repeats"
                     /note="LRRCT; Leucine rich repeat C-terminal domain.
                     Leucine Rich Repeats pfam00560 are short sequence motifs
                     present in a number of proteins with diverse functions and
                     cellular locations. Leucine Rich Repeats are often flanked
                     by cysteine rich domains. This domain is often found at
                     the C-terminus of tandem leucine rich repeats"
                     /db_xref="CDD:pfam01463"
     CDS             1..347
                     /gene="LRG"
                     /coded_by="NM_052972.1:12..1055"
                     /note="contains leucine-rich repeat"
                     /db_xref="GeneID:116844"
ORIGIN
        1 msswsrqrpk spggiqphvs rtlfllllla asawgvtlsp kdcqvfrsdh gssiscqppa
       61 eipgylpadt vhlaveffnl thlpanllqg asklqelhls snglesls pe flrpvpqlrv
      121 ldltrnaltg lppglfqasa tldtlvlken qlevlevswl hglkalghld lsgnrlrklp
      181 pgllanftll rtldlgenql etlppdllrg plqlerlhle gnklqvlgkd lllpqpdlry
      241 lflngnklar vaagafqglr qldmldlsnn slasvpeglw aslgqpnwdm rdgfdisgnp
      301 wicdqnlsdl yrwlqaqkdk mfsqndtrca gpeavkgqtl lavaksq
//
```